

Sulphur and iron abundances in halo stars¹

Poul E. Nissen

Department of Physics and Astronomy, University of Aarhus, Denmark

YuQin Chen

National Astronomical Observatories, Beijing, China

Martin Asplund

Mount Stromlo Observatory, Australian National University

Max Pettini

Institute of Astronomy, Cambridge, UK

Abstract.

From equivalent widths of the S I lines at 8694 Å, Israelian & Rebolo (2001) and Takada-Hidai et al. (2002) have derived a surprisingly high sulphur-to-iron ratio ($[S/Fe] \simeq 0.5$ to 0.7) in six halo stars with $[Fe/H] \simeq -2.0$ suggesting perhaps that hypernovae made a significant contribution to the formation of elements in the early Galaxy. To investigate this problem we have used high-resolution spectra obtained with the ESO VLT/UVES spectrograph to determine the S/Fe ratio in 19 main-sequence and subgiant stars ranging in $[Fe/H]$ from -3.2 to -0.7 . The sulphur abundances are determined from S I lines at 8694 Å and 9212 - 9237 Å, and the iron abundances from about 20 Fe II lines. S/Fe ratios as derived from 1D model atmospheres are presented and possible 3D effects are discussed. The initial results from our survey do not confirm the high values of $[S/Fe]$ quoted above; instead we find that the ratio $[S/Fe]$ remains constant at about 0.35 dex for metallicities $-3 < [Fe/H] < -1$.

1. Introduction

Sulphur is generally regarded as an α -capture element. The work on Galactic stars by François (1987, 1988) supported this view by showing that $[S/Fe]$ increases from zero at solar metallicities to a plateau level of about +0.5 dex in the metallicity range $-1.8 < [Fe/H] < -0.8$, an analogous behaviour to that of other α -elements Mg, Si, and Ca, see Ryan et al. (1996). The standard inter-

¹Based on observations collected at the European Southern Observatory, Chile (ESO No. 67.D-0106)

pretation is that this trend arises from the time delay in the production of $2/3$ of the iron by SN of Type Ia relative to the near-instantaneous release of the α -elements by Type II SN.

Recent observations of sulphur in metal-poor stars by Israelian & Rebolo (2001) have, however, challenged this view. Their data suggest that $[S/Fe]$ increases linearly with decreasing $[Fe/H]$ to a level as high as $[S/Fe] \sim +0.7$ at $[Fe/H] = -2.0$. The study of Takada-Hidai et al. (2002) based on Keck HIRES observations supports a quasi linear dependence of $[S/Fe]$ on $[Fe/H]$ although in their case $[S/Fe]$ reaches only 0.5 dex at $[Fe/H] = -2.0$.

As a possible explanation of the high value of $[S/Fe]$ in metal-poor stars, Israelian & Rebolo propose that massive supernovae with exploding He-cores and a high explosion energy make a significant contribution to the early chemical evolution of galaxies. According to Nakamura et al. (2001) these hypernovae overproduce S with respect to O, Mg and Fe. With this intriguing possibility in mind a more thorough investigation of sulphur abundances in halo stars seems worthwhile. A clarification of the trend of S abundances is also much needed in deciphering the chemical enrichment of damped Ly α systems (DLAs), widely regarded as the progenitors of present-day galaxies at high redshift. Its importance stems from the fact that, unlike most other heavy elements, S is not depleted onto dust. Consequently, observations of the relatively weak S II triplet resonance lines at $\lambda\lambda 1250, 1253, 1259$ yield a direct measurement of the abundance of S in DLAs. The only other element for which this is the case is Zn, and indeed most of our current knowledge of the chemical evolution of the universe at high redshift is based on surveys of $[Zn/H]$ in DLAs (e.g. Pettini et al. 1999, Prochaska & Wolfe 2002). If S is an α -capture element, then its abundance relative to Zn (an iron-peak element) could be used as ‘a chemical clock’ to date the star-formation process at high z . Specifically, if the major star formation episodes in the DLAs observed occurred within the last ≈ 0.5 Gyr, we would expect to measure enhanced $[S/Zn]$ ratios, and *vice versa*. Data on the abundance of S in DLAs have been relatively scarce until recently, but are now becoming available at a progressively faster rate thanks largely to the high sensitivity of the VLT/UVES high resolution spectrograph at blue and near-UV wavelengths. The picture is still confused (Centuri n et al. 2000, Prochaska & Wolfe 2002), but one thing is clear: without a secure knowledge of the behaviour of S in metal-poor Galactic stars we stand no hope of interpreting the situation at high z .

In this paper we report on a large scale survey of sulphur abundances in metal-poor stars carried out with VLT/UVES. Effects on the derived S/Fe ratio from the modelling of stellar atmospheres are discussed, and preliminary results for about half of the 35 stars observed are presented.

2. Observations and data reduction

Previous determinations of sulphur abundances in metal-poor stars, including those of Israelian & Rebolo (2001) and Takada-Hidai et al. (2002), have been based on the high excitation ($\chi = 7.87$ eV) $\lambda\lambda 8693.9, 8694.6$ S I lines. They are, however, very weak in metal-poor dwarfs and giants ($W < 3$ mÅ at $[Fe/H] \sim -2.0$), and practically impossible to observe at $[Fe/H] \sim -2.5$. In our survey




Figure 1. The VLT/UVES spectrum of HD 110621 ($[\text{Fe}/\text{H}] = -1.66$). The dotted line shows the spectrum before removal of the numerous telluric H_2O lines. The thick, full drawn line is the spectrum after division with the spectrum of the B-type star HR 5488 using the IRAF task ‘*telluric*’ to obtain the best fit between the two sets of H_2O lines. The radial velocity of HD 110621 was 207.1 km s^{-1} , so all stellar lines are shifted by about 6.4 Å . The Si I triplet, a Mg II line and the Paschen ζ H I line of HD 110621 are seen. The broad ‘emission’ feature is due to the H I line in the B star spectrum. Excessive noise in the stellar spectrum are seen where strong H_2O lines have been removed.

we have taken advantage of the high efficiency of the VLT/UVES instrument and concentrated on the stronger Si I triplet ($\chi = 6.52 \text{ eV}$) at $9212 - 9238 \text{ Å}$. The dichroic mode of UVES was used to cover the spectral region $6700 - 10000 \text{ Å}$ in the red arm and $4000 - 5000 \text{ Å}$ in the blue arm; the latter includes a number of Fe II lines suitable for determining the iron abundance. The resolution of the spectra is $\lambda/\Delta\lambda \simeq 60\,000$ with 4 pixels per spectral resolution element. Typical S/N ratios are 300 in the blue region and 200 at the $9212 - 9238 \text{ Å}$ triplet.

The red spectra were reduced using the echelle package in IRAF. A problem with the $9212 - 9238 \text{ Å}$ region is the presence of numerous, quite strong telluric H_2O lines, see Fig. 1. To remove these lines fast rotating, early-type stars were observed on each night and reduced in the same way as the program stars. The IRAF task ‘*telluric*’ was then used to remove the telluric lines. When the early-type star is observed at about the same airmass as the program star this technique also serves to remove a residual fringing remaining from the flatfielding of the spectra.




Figure 2. A sequence of spectra around the $\lambda\lambda 8693.9, 8694.6$ Si I lines.




Figure 3. A sequence of spectra around the $\lambda 9212.9$ Si I line. The arrows indicate positions where telluric H_2O lines have been removed. The spectra have S/N around 200.

Representative spectra of stars are shown in Fig. 2 (8694 Å region) and Fig. 3 (9212 Å region). As seen the Si I line at 9212.9 Å can be clearly detected in a $[\text{Fe}/\text{H}] \sim -3.2$ star. Eight stars were observed twice on different nights; a comparison shows that the equivalent widths of the $\lambda\lambda 8693.9, 8694.6$ lines are measured with an accuracy of about $\pm 0.5 \text{ mÅ}$ whereas the $\lambda\lambda 9212.9, 9237.5$ lines are measured to an accuracy of $\pm 2 \text{ mÅ}$.

The blue spectra were reduced with MIDAS routines. For a few stars it was checked that IRAF reduction gives practically the same result. These blue spectra have $S/N \sim 300$ and allowed us to measure accurate equivalent widths of about 20 weak Fe II lines.

3. Modelling of the stellar atmospheres

The measured equivalent widths were used to derive S and Fe abundances based on 1D MARCS (Asplund et al. 1997) model atmospheres assuming LTE. Effective temperatures were derived from the color indices $b-y$ and $V-K$, and surface gravities via Hipparcos parallaxes and/or the Strömgren c_1 index. For details we refer to the recent paper by Nissen et al. (2002) on oxygen abundances.

Oscillator strengths of the blue Fe II lines were determined from an inverted abundance analysis of three stars (HD 103723, HD 160617 and HD 140283) adopting $[\text{Fe}/\text{H}]$ values determined by Nissen et al. (2002) from very accurate equivalent widths of weak Fe II lines analyzed differentially with respect to the solar spectrum. This ensures that we are on the same $[\text{Fe}/\text{H}]$ scale as Nissen et al. (2002).

Oscillator strengths of the sulphur lines were taken from Lambert & Luck (1978). We note that the values for the 8693.9 and 8694.6 Å lines ($\log gf = -0.56$ and 0.03 , respectively) and the solar spectrum equivalent widths provide a sulphur abundance of the Sun very close to the meteoritic value, $\log \epsilon(\text{S}) = 7.20$. The 9212.9 and 9237.5 Å Si I are too strong in the solar spectrum to provide reliable abundances, but we note that the sulphur abundance derived from this pair of lines agree well with that derived from the $\lambda\lambda 8693.9, 8694.6$ pair for 12 stars in the metallicity range $-1.5 < [\text{Fe}/\text{H}] < -0.7$. The mean difference between the S abundance derived from the $\lambda\lambda 8693.9, 8694.6$ pair and the 9212.9, 9237.5 Å lines is $+0.03$ dex with an rms deviation of the difference of ± 0.10 dex. The third line of the near-infrared Si I triplet at 9228.1 Å is unfortunately very close to the Paschen ζ H I line at 9229.0 Å and could not be used for a reliable determination of the sulphur abundance.

Errors in T_{eff} ($\pm 70 \text{ K}$) and $\log g$ (± 0.15 dex) have only a minor effect on the derived $[\text{S}/\text{Fe}]$ (less than 0.05 dex). This is connected to the fact that the abundances of both elements are determined from their main ionization stages, Si I and Fe II, respectively. The microturbulence was determined from requiring that the abundance derived from the 20 Fe II lines should be independent of W , or in the case of stars with very weak Fe II lines ($[\text{Fe}/\text{H}] < -1.5$) ξ_{micro} is assumed to be 1.5 km s^{-1} . In any case, the effect of the uncertainty of ξ_{micro} on the S abundance is quite negligible due to the weakness of the Si I lines.

As shown by Takada-Hidai et al. (2002) non-LTE effects on the weak, high excitation Si I lines are likely to be small (< 0.05 dex). The same is the case for the Fe II lines according to Thévenin & Idiart (1999).

Table 1. 1D - 3D model atmosphere effects on sulphur abundances. For each set of model atmosphere parameters the table first gives the assumed 1D sulphur abundances ($A = \log \epsilon(\text{S})$), then the computed 1D equivalent widths for the three most important Si I lines and the 3D abundances derived from these equivalent widths.

T_{eff}	$\log g$	[Fe/H]	8694.6 Å			9212.8 Å			9237.5 Å		
			A_{1D}	$W_{1D}(\text{mÅ})$	A_{3D}	$W_{1D}(\text{mÅ})$	A_{3D}	$W_{1D}(\text{mÅ})$	A_{3D}	$W_{1D}(\text{mÅ})$	A_{3D}
5767	4.44	0.0	7.33	35.7	7.29	143.5	7.32	109.3	7.32		
5822	4.44	-1.0	6.73	15.2	6.75	112.4	6.75	81.2	6.75		
5837	4.44	-2.0	5.73	1.7	5.79	42.5	5.86	23.3	5.84		
5890	4.44	-3.0	4.73	0.2	4.78	7.1	4.81	3.1	4.81		
6191	4.04	0.0	7.33	55.6	7.36	160.9	7.44	127.3	7.42		
6180	4.04	-1.0	6.73	25.8	6.77	114.0	6.74	88.1	6.76		
6178	4.04	-2.0	5.73	3.5	5.77	51.9	5.82	32.7	5.82		
6205	4.04	-3.0	4.73	0.4	4.77	11.4	4.81	5.2	4.80		

Much attention has recently been paid to the effect on abundance determinations of 3D hydrodynamical modelling of convection and granulation in stellar atmospheres. As first shown by Asplund et al. (1999) there is a very significant surface cooling in metal-poor stars caused by the adiabatic cooling of rising and expanding elements coupled with the lack of radiative heating due to the low line opacity in metal-poor stars. The effect on lines formed in the upper atmosphere can be very large. As an example Asplund & García Pérez (2001) estimated 3D effects on oxygen abundances derived from OH lines of the order of -0.6 dex for turnoff stars with $[\text{Fe}/\text{H}] < -2.0$.

To investigate the 3D effects on the sulphur lines used in the present study a differential 1D - 3D study similar to that described by Asplund & García Pérez (2001) has been carried out. The results are given in Table 1 for two sets of model atmospheres with metallicities of $[\text{Fe}/\text{H}] = 0.0, -1.0, -2.0$ and -3.0 . The first set has T_{eff} and $\log g$ close to the solar values, whereas the second set represents stars close to the turnoff. For each model atmosphere Table 1 first gives the adopted S abundance, then the corresponding computed 1D equivalent width, and finally the S abundance calculated from the 1D equivalent width on the basis of the 3D model atmosphere. As seen the maximum 3D effect occurs for the most metal-poor stars: the 3D abundance is increased by 0.07 to 0.13 dex with respect to the 1D abundance in the case of the $\lambda\lambda 9212.9, 9237.5$ Si I lines. Interestingly, there is, however, a similar 3D effect ($\sim +0.10$ dex) on the Fe abundance derived from Fe II lines (Nissen et al. 2002). Hence, it appears that we are in the favourable situation that the S/Fe ratio is quite immune to 1D - 3D effects.

4. Results

Up to now we have reduced and analyzed about half of the VLT/UVES spectra. The derived sulphur abundances for 19 halo stars are given in Table 2 together

with atmospheric parameters and iron abundances. Fig. 4 shows a plot of $[\text{S}/\text{Fe}]$ vs. $[\text{Fe}/\text{H}]$ with disk stars from Chen et al. (2002) added. The sulphur abundances of the disk stars were derived from the $\lambda\lambda 8693.9, 8694.6$ S I lines and three other weak S I lines in the spectral range 6000 – 6800 Å.

Table 2. Effective temperatures, surface gravities, iron and sulphur abundances for 19 stars with halo kinematics.

Star	T_{eff}	$\log g$	$[\text{Fe}/\text{H}]$	$[\text{S}/\text{H}]$	$[\text{S}/\text{Fe}]$	Note
HD103723	6034	4.26	−0.81	−0.84	−0.03	α -def.
HD105004	5919	4.36	−0.86	−0.85	0.01	α -def.
HD106038	5919	4.30	−1.42	−1.00	0.42	
HD108177	6039	4.25	−1.74	−1.39	0.35	
HD110621	5981	3.99	−1.66	−1.34	0.32	
HD121004	5636	4.33	−0.76	−0.45	0.31	
HD140283	5692	3.69	−2.41	−2.11	0.30	
HD146296	5666	4.18	−0.72	−0.54	0.18	
HD148816	5828	4.14	−0.73	−0.50	0.23	
HD160617	5931	3.77	−1.80	−1.40	0.40	
G11-44	5994	4.29	−2.08	−1.71	0.37	
G13-09	6361	4.00	−2.27	−1.81	0.46	
G16-13	5603	4.17	−0.75	−0.47	0.28	
G20-08	5855	4.16	−2.27	−1.86	0.41	
G53-41	5829	4.16	−1.34	−1.00	0.34	
G64-12	6512	4.39	−3.16	−2.80	0.36	
G66-30	6345	4.24	−1.52	−1.30	0.22	
BD−13 3442	6501	4.18	−2.59	−2.30	0.29	
CD−42 14278	5804	4.25	−2.13	−1.82	0.31	

As seen from Fig. 4 $[\text{S}/\text{Fe}]$ shows a clear plateau ($[\text{S}/\text{Fe}] \simeq 0.35$) for $[\text{Fe}/\text{H}]$ below -1.2 . There is no tendency for a linear rise of $[\text{S}/\text{Fe}]$ as claimed by Israelian & Rebolo (2001) and Takada-Hidai et al. (2002). We note in this connection that their samples include only three stars with $[\text{Fe}/\text{H}] < -2.0$ and none below $[\text{Fe}/\text{H}] = -2.3$, whereas our sample includes seven stars in the range $-3.2 < [\text{Fe}/\text{H}] < -2.0$. Furthermore, the error bars on their sulphur abundance determinations are large due to the weakness of the S I 8694.6 Å line; the 9212.8 and 9237.5 Å S I lines, on the other hand, are rather ideal for sulphur abundance determinations below $[\text{Fe}/\text{H}] = -2.0$. Although we cannot exclude the existence of a few stars with high values of S/Fe^2 it would appear that sulphur shows the same trend as other typical α -capture elements Mg, Si and Ca. Hence, our preliminary conclusion is that there is no need to invoke element production by hypernovae or very massive supernovae to explain the general behaviour of sulphur. Traditional Galactic evolution models with near-instantaneous production of α -elements by Type II supernovae and delayed production of the iron-peak

²We do confirm the high value $[\text{S}/\text{Fe}] \simeq 0.7$ of HD 19445 ($[\text{Fe}/\text{H}] \simeq -1.9$) if we adopt the equivalent width (2.7 mÅ) of the S I 8694.6 Å line given by Israelian & Rebolo based on their very high S/N spectrum




Figure 4. $[S/Fe]$ vs. $[Fe/H]$. Filled circles are halo stars from the present work and open circles are disk stars from Chen et al. (2002). The typical error bars on the abundance ratios are shown, and the two α -deficient halo stars, HD 103723 and HD 105004 (Nissen & Schuster 1997), are indicated.

elements are in good agreement with our data for sulphur (Chiappini et al. 1999, Goswami & Prantzos 2000).

An interesting detail in Fig. 4 should be noted. All stars from the present investigation (plotted with filled circles) have halo kinematics, including six stars with $-0.9 < [Fe/H] < -0.7$. They have Galactic rotational velocities less than 50 km s^{-1} i.e. well below the characteristic rotational velocity of disk stars (225 km s^{-1}) and thick disk stars (175 km s^{-1}). As seen from the figure, there is then an overlap between halo stars and thick disk stars in the metallicity range $-1.0 < [Fe/H] < -0.6$. Out of the six stars four have enhanced S/Fe ratios like the thick disk stars but two (HD 103723 and HD 105004) have a solar S/Fe ratio. Interestingly, HD 103723 and HD 105004 were also found to be deficient in other α -element (O, Mg, Si, Ca, Ti) by Nissen & Schuster (1997). On the basis of their Galactic orbits, these authors speculated that the α -deficient stars have been accreted from dwarf galaxies with a slow chemical evolution in contrast to the ‘normal’ halo stars with enhanced $[\alpha/Fe]$ formed in a fast chemical evolution in the inner part of our Galaxy. The fact that S and the classical α -elements, Mg, Si and Ca, are deficient in the same stars is a further indication that sulphur belongs to this group of elements. Thus there seems to be little ground, on the basis of our data, for the reservations expressed by Prochaska et al. (2000) concerning use of S as an α -element in the analysis of abundance ratios in DLAs.

On the other hand, one should remain aware that the overabundance of the α -elements at low metallicities need not be a universal phenomenon. The finding by Nissen & Schuster (1997) that even some stars in the Milky Way do not fit this pattern is a warning against the naive assumption that such an overabundance should also apply to all metal-poor DLAs.

References

- Asplund M., Gustafsson B., Kiselman D., Eriksson K. 1997, *A&A*, 318, 521
Asplund M., Nordlund Å., Trampedach R., Stein R.F. 1999, *A&A*, 346, L17
Asplund M., García Pérez A.E. 2001, *A&A*, 372, 601
Centurión M., Bonifacio P., Molaro P., Vladilo G. 2000, *ApJ*, 536, 540
Chen Y.Q., Nissen P.E., Zhao G., Asplund M. 2002, *A&A* in press, (astro-ph/0206075)
Chiappini C., Matteucci F., Beers T.C., Nomoto K. 1999, *ApJ*, 515, 226
François P. 1987, *A&A*, 176, 294
François P. 1988, *A&A*, 195, 226
Goswami A., Prantzos N. 2000, *A&A*, 359, 191
Israelian G., Rebolo R. 2001, *ApJ*, 557, L43
Lambert D.L., Luck R.E. 1978, *MNRAS*, 183, 79
Nissen P.E., Schuster W.J. 1997, *A&A*, 326, 751
Nissen P.E., Primas F., Asplund M., Lambert D.L. 2002, *A&A* in press, (astro-ph/0205372)
Pettini M., Ellison S.L., Steidel C.C., Bowen D.V. 1999, *ApJ*, 510, 576
Prochaska J. X., Naumov S.O., Carney B.W. et al. 2000, *AJ*, 120, 2513
Prochaska J. X., Wolfe A.M. 2002, *ApJ*, 566, 68
Ryan S.G., Norris J.E., Beers T.C. 1996, *ApJ*, 471, 254
Takada-Hidai M., Takeda Y., Sato S., et al. 2002, *ApJ*, 573, 584
Thévenin F., Idiart T.P. 1999, *ApJ*, 521, 753
Nakamura T., Umeda H., Iwamoto K.I. et al. 2001, *ApJ*, 555, 880